


Diagnosing Circumstellar Debris Disks

Joseph M. Hahn
Space Science Institute



the edge-on debris disk orbiting β Pictoris, from Heap et al (2000)

Dusty circumstellar debris disks:


sites of ongoing planet formation?
or planetesimal destruction?

the optimistic view:

- dust lifetimes \ll host star's age
- requires replenishment, presumably by unseen planetesimals
- planetesimals are the seeds of planets

but models of planetesimal collisional/accretional evolution
in outer Solar System show (Stern & Colwell 1997, Kenyon 2002):

- planet formation in the $r \gtrsim 30$ AU zone is very inefficient, requiring $M_p \sim 30 M_\oplus$ just to form a handful of Plutos
- much of the leftover mass then grinds down to dust, blown away by radiation pressure (RP) in $t \sim 500$ Myrs
- characteristic dust mass loss rate is $\dot{M}_d \sim M_p/t \sim 10^{13}$ gm/sec




AU Mic, from Fitzgerald et al 2007

Do disk observations support this finding...

that planet-formation is lossy and inefficient at $r \gtrsim 30$ AU?

To address this,

- develop a model of a debris disk
- fit to observations
- hopefully say something about the disk's prospects for planet formation




AU Mic surface brightness
from Strubbe & Chiang (2006)

The relevant physics is described in Strubbe & Chiang (2006),

- unseen planetesimals collide & generate dust
- RP lofts smaller grains into wide orbits, $r \sim 100$'s of AU
- collisions among dust shatter grains until $R < R_{\text{blowout}}$

The model:

- quantize the problem, so $\int \rightarrow \sum$
- $1 \leq N_r \leq 5$ circular planetesimal rings that produce dust at $N_\ell = 100$ longitudes
- dust have $N_R = 200$ dust size-bins
- dust production rate is power-law in size, $d\dot{N}(R) \propto R^{-q}$



Dust grains have size parameter $\beta = \frac{RP}{G} \sim 0.6/R_{\mu\text{m}}$ (if star is solar),

and dust orbit elements are simple functions of β (S&C2006):

$$a(\beta) = \frac{1 - \beta}{1 - 2\beta} r_p \quad \text{and} \quad e(\beta) = \frac{\beta}{1 - \beta}$$

so bound dust have $\beta < \frac{1}{2}$ and radii $R > R_{\text{blowout}}$

where $R_{\text{blowout}} \sim 1 \mu\text{m}$ (when solar)

Dust abundance obeys rate equation:

$N_i(t)$ = no. of grains of radius R_i in orbit $a_i, e_i, \tilde{\omega}_i$

$$\frac{dN_i}{dt} = P_i - \sum_j \alpha_{ij} N_i N_j$$

= production - destruction


which is solved numerically for $N_i(t)$

α_{ij} = probability per time for a grain in orbit i to collide with grain in orbit j
 = function($a_i, a_j, e_i, e_j, \tilde{\omega}_i, \tilde{\omega}_j, R_i, R_j$)

Impact must also be fast enough for grain j to shatter grain i :

$$|\mathbf{v}_j - \mathbf{v}_i|^2 \gtrsim Q^* \left(\frac{R_i}{R_j} \right)^3$$

where Q^* is dust strength



Example:

$r_p = 50$ AU p'mal ring

$\dot{M}_d = 10^{13}$ gm/sec

$Q^* = 10^6$ ergs/gm (weak)

$I = 0.1$ rad = 6°

rate equation

provides scale-factors:


abundance



$$N_0 \propto r_p^{7/4} \sqrt{I \dot{M}_d}$$

timescale

$$T_0 \propto r_p^{7/4} \sqrt{I / \dot{M}_d}$$

⇒ heavier dust production results in a more massive debris disk that settles faster into collisional equilibrium





Dust collisional lifetimes: $T_c(R) = \frac{N(R)}{P(R)}$



when dust grains are weak, $Q^* < 10^6$ ergs/gm, all collisions are destructive,

$T_c \propto \dot{M}_d^{-1/2}$ and $T_c \propto R^{-2}$ for $R \gtrsim 2R_{\text{blowout}}$

\Rightarrow large grains have **short** lifetimes due to bombardment by abundant small grains

increasing Q^* increases lifetime of large dust that are confined to planetesimal disk

Disk optical depth τ



Surface brightness (SB) of edge-on disk

β Pic, AU Mic are seen edge-on


their SB is sensitive to
asymmetry in light scattering

$$g = \int \Phi(\phi) \cos \phi d\Omega$$

when $g = 0$ (isotropic scattering)
inner SB($x < r_{in}$) is flat if
planetesimal disk has donut-hole

if $|g| \gtrsim 0.7$ (forward scattering),
then SB has a knee-bend
where LOS passes thru
planetesimal disk

where $SB(x) \propto x^{-7/2}$
indicates planetesimal r_{out}




planetesimals reside at $r_{in} < r < r_{out}$
where $r_{in} = 0.5r_{out}$

Diagnosing β Pictoris

fit requires:

- broad planetesimal disk,
 $75 \lesssim r_p \lesssim 150$ AU
- heavy dust production
 $\dot{M}_d \sim 3 \times 10^{15}$ gm/sec
(300 \times higher than S&C model)
- grains are probably reflective.
I assumed $Q_s = 0.7$,
similar to Saturn's icy rings
 - note $SB \propto Q_s \sqrt{\dot{M}_d}$,
if $Q_s = 0.1$ (dark dust)
then $\dot{M}_d \uparrow \times 100$
- dust size dist' has $q = 2.5$,
shallower than Dohnanyi $q = 3.5$
- dust grains are strong,
 $Q^* \sim 10^8$ ergs/gm,
to preserve large grains at $x \sim 100$ AU

optical HST observations by
Golimowski et al (2006)




- knee indicates that dust are
asymmetric light scatters, $g \simeq 0.7$

Mass of β Pic Disk

Assuming Bond albedo $Q_s = 0.7$:

- $M_{dust} \simeq 11$ lunar masses
 - comparable to estimate from sub-mm observations by Holland et al (1998)
- dust cross section is $A_{dust} \simeq 2 \times 10^{20} \text{ km}^2$
- note star's age $t_\star \simeq 12$ Myrs, so implied mass-loss is $\dot{M}_d t_\star \sim 160 M_\oplus!$
 - β Pic's planetesimal disk is (or was) very massive!




inferred optical depth $\tau(r)$
assuming dark $Q_s = 0.1$ and
bright $Q_s = 1$ grains

The prospects for planet formation at β Pic are...

...unclear? grim?

- the planetesimal disk is suffering heavy mass loss due to collisional grinding + blowout by RP,
 $\dot{M}_p \sim 13 M_{\oplus}/\text{Myr}$.
- β Pic's planetesimal disk is or was very massive,
 $M_p \gtrsim 160 M_{\oplus}$ in $75 \lesssim r \lesssim 150$ AU zone
- I suspect that the $r \gtrsim 75$ AU zone at β Pic may be a region of planetesimal destruction, rather than a site of future planet formation



β Pic with radial variations factored out

Next steps

- I also need to model the disk's thermal emission
 - fits to optical + sub-mm observations will allow me to pin down \dot{M}_d and Q_s with greater certainty
- will couple this debris-disk model to Stu W's planetesimal model
 - his code can track the growth and erosion of planetesimals
 - this will produce a more realistic treatment of the disk's dust production rate $\dot{M}_d(t)$ over time
 - will also allow us to infer or else constrain the unseen planetesimal disk mass with greater realism
- preprint will be available
- supported by Hubble Theory/Archive research program